



Pisani, D., Pett, W., Dohrmann, M., Feuda, R., Rota-Stabelli, O., Philippe, H., Lartillot, N., & Wörheide, G. (2016). Reply to Halanych et al. Ctenophore misplacement is corroborated by independent datasets. *Proceedings of the National Academy of Sciences of the United States of America*, 113(8), E948-949.  
<https://doi.org/10.1073/pnas.1525718113>

Peer reviewed version

License (if available):  
Unspecified

Link to published version (if available):  
[10.1073/pnas.1525718113](https://doi.org/10.1073/pnas.1525718113)

[Link to publication record in Explore Bristol Research](#)  
PDF-document

## University of Bristol - Explore Bristol Research

### General rights

This document is made available in accordance with publisher policies. Please cite only the published version using the reference above. Full terms of use are available:  
<http://www.bristol.ac.uk/red/research-policy/pure/user-guides/ebr-terms/>

Letter: Biological Sciences: Evolution

**Reply to Halanych *et al.*: Ctenophore misplacement is corroborated by independent datasets.**

Davide Pisani<sup>a,b,1\*</sup>, Walker Pett<sup>c\*</sup>, Martin Dohrmann<sup>d</sup>, Roberto Feuda<sup>e</sup>, Omar Rota-Stabelli<sup>f</sup>,  
Hervé Philippe<sup>g,h</sup>, Nicolas Lartillot<sup>c</sup>, Gert Wörheide<sup>d,i,1</sup>

*<sup>a</sup>School of Earth Sciences, University of Bristol, Bristol BS8 1TG, United Kingdom; <sup>b</sup>School of Biological Sciences, University of Bristol, Bristol BS8 1TG, United Kingdom; <sup>c</sup>Laboratoire de Biométrie et Biologie Évolutive, Université Lyon 1, CNRS, UMR 5558, 69622 Villeurbanne cedex, France; <sup>d</sup>Department of Earth & Environmental Sciences & GeoBio-Center, Ludwig-Maximilians-Universität München, Munich 80333, Germany; <sup>e</sup>Division of Biology and Biological Engineering, California Institute of Technology, Pasadena, CA 91125; <sup>f</sup>Department of Sustainable Agro-Ecosystems and Bioresources, Research and Innovation Centre, Fondazione Edmund Mach, San Michele all' Adige 38010, Italy; <sup>g</sup>Centre for Biodiversity Theory and Modelling, USR CNRS 2936, Station d'Ecologie Expérimentale du CNRS, Moulis 09200, France; <sup>h</sup>Département de Biochimie, Centre Robert-Cedergren, Université de Montréal, Montreal, QC, Canada H3C 3J7; and <sup>i</sup>Bayerische Staatssammlung für Paläontologie und Geologie, Munich 80333, Germany.*

1: Corresponding authors: Davide Pisani ([davide.pisani@bristol.ac.uk](mailto:davide.pisani@bristol.ac.uk)) and Gert Wörheide ([woerheide@lmu.de](mailto:woerheide@lmu.de)).

\*: These authors equally contributed to the work: Dual first authorship.



In their letter, Halanych *et al.* (1) criticize our recent assertion (2) that the phylogenetic placement of ctenophores as the sister group to all other animals (the Ctenophora-sister hypothesis) in three previous studies (3–5) was an artifact caused by undetected systematic error.

Halanych *et al.* claim we used no “objective criteria” to identify sources of systematic error. In fact, we used an objective comparison of Bayesian cross-validation scores to select the best-fitting substitution model, since poorly-fitting models are a frequent source of systematic error. Halanych *et al.* point out that this comparison did not include partitioned site-homogeneous models. However, they did not mention that only one of the studies we addressed (3) used this approach, and that multiple site-homogeneous partitions still do not account for within-partition site-heterogeneous biochemical constraints, which our results showed had a major impact on model fit and the tree topology.

Halanych *et al.* also incorrectly suggest that our model-selection procedure relied on circular assumptions about the position of sponges. No such assumptions were made to demonstrate the better fit of site-heterogeneous models. In fact, our cross-validation tests were conservative in favor of Ctenophora-sister, because we “trained [all models] under the tree topology favored by WAG [Ctenophora-sister], thus making the test conservative in favor of the WAG model [and consequently Ctenophora-sister]” (Methods in (2)).

Halanych *et al.* further claim our conclusions rested extensively on unconverged analyses. In reality, our conclusions about the position of Ctenophora were based solely on converged analyses, with two unconverged analyses mentioned only for completeness in one case, or to demonstrate weak phylogenetic signal in the other (reanalysis of (5)).

The main criticism of Halanych *et al.* is that we arbitrarily preferred results obtained using only closely-related outgroups. In fact, our results simply showed weaker average support

for Ctenophora-sister under better-fitting models. When only holozoan (not just choanoflagellate) outgroups were included, support for Ctenophora-sister did deteriorate (Figure 2B-C in (2)), or support for Porifera-sister emerged (Figure 2A in (2)), and this effect became more pronounced when using only choanoflagellates. This sensitivity of the position of Ctenophora to model choice after excluding distant outgroups suggests these outgroups exacerbate systematic error by obscuring the effect of better-fitting models.

The final comment (1) about the questionable use of gene content data is unjustified. The observation that “gene content varies greatly within phyla” is evidence that it contains phylogenetic signal. Accordingly, analysis of gene content (corrected for ascertainment bias) recovered well-established bilaterian relationships (Figure 3 in (2)), and corroborated non-bilaterian relationships inferred from amino acid data (Figure 1 in (2)).

Finally, Halanych *et al.* claim that our conclusions mistreated complex traits as single characters and ignored the position of Placozoa. In fact, we acknowledged that “Future studies [...] will help to clarify the relationship between the homology of similar structures and their underlying genetic mechanisms”, and “resolving the exact phylogenetic positions of Ctenophora and Placozoa [...] will be crucial to reconstruct the evolution of key characters, such as nervous systems, muscles, and digestive tracts, in more detail” (2).

## References

1. Halanych KM, Whelan NV, Kocot KM, Kohn AB, Moroz LL (2016) Miscues misplace sponges. *Proc Natl Acad Sci*.
2. Pisani D, *et al.* (2015) Genomic data do not support comb jellies as the sister group to all other animals. *Proc Natl Acad Sci* 112(50):15402–15407.
3. Whelan NV, Kocot KM, Moroz LL, Halanych KM (2015) Error, signal, and the placement of Ctenophora sister to all other animals. *Proc Natl Acad Sci* 112(18):5773–5778.

4. Ryan JF, *et al.* (2013) The genome of the ctenophore *Mnemiopsis leidyi* and its implications for cell type evolution. *Science* 342(6164):1242592.
5. Moroz LL, *et al.* (2014) The ctenophore genome and the evolutionary origins of neural systems. *Nature* 510(7503):109–114.